


**TEST PAPER OF JEE(MAIN) EXAMINATION – 2019**
**(Held On Wednesday 09<sup>th</sup> JANUARY, 2019) TIME : 2 : 30 PM To 05 : 30 PM**
**CHEMISTRY**


1. Good reducing nature of  $H_3PO_2$  attributed to the presence of:

- (1) One P-OH bond      (2) One P-H bond  
 (3) Two P-H bonds      (4) Two P-OH bonds

Ans. (3)

Sol.  $H_3PO_2$  is good reducing agent due to presence

of two P-H bonds.



2. The complex that has highest crystal field splitting energy ( $\Delta$ ), is :

- (1)  $K_3[Co(CN)_6]$   
 (2)  $[Co(NH_3)_5(H_2O)]Cl_3$   
 (3)  $K_2[CoCl_4]$   
 (4)  $[Co(NH_3)_5Cl]Cl_2$

Ans. (1)


Sol. As complex  $K_3[Co(CN)_6]$  have  $CN^-$  ligand which is strong field ligand amongst the given ligands in other complexes.

3. The metal that forms nitride by reacting directly with  $N_2$  of air, is :

- (1) K      (2) Cs      (3) Li      (4) Rb

Ans. (3)

Sol. Only Li react directly with  $N_2$  out of alkali metals



4. In which of the following processes, the bond order has increased and paramagnetic character has changed to diamagnetic ?






- (1)  $N_2 \rightarrow N_2^+$       (2)  $NO \rightarrow NO^+$   
 (3)  $O_2 \rightarrow O_2^{2-}$       (4)  $O_2 \rightarrow O_2^+$

Ans. (2)

Sol.


Process	Change in magnetic nature	Bond Order Change
$N_2 \rightarrow N_2^+$	Dia $\rightarrow$ para	3 $\rightarrow$ 2.5
$NO \rightarrow NO^+$	Para $\rightarrow$ Dia	2.5 $\rightarrow$ 3
$O_2 \rightarrow O_2^{2-}$	Para $\rightarrow$ Dia	2 $\rightarrow$ 1
$O_2 \rightarrow O_2^+$	Para $\rightarrow$ Para	2 $\rightarrow$ 2.5

5. The major product of the following reaction is:



Ans. (4)

Sol.



**JEE (Main) Examination–2019/Evening Session/09-01-2019**

6. The transition element that has lowest enthalpy of atomisation, is :

- (1) Zn
- (2) Cu
- (3) V
- (4) Fe

Ans. (2)

Sol. Since Zn is not a transition element so transition element having lowest atomisation energy out of Cu, V, Fe is Cu.

7. Which of the following combination of statements is true regarding the interpretation of the atomic orbitals ?

- (a) An electron in an orbital of high angular momentum stays away from the nucleus than an electron in the orbital of lower angular momentum.
- (b) For a given value of the principal quantum number, the size of the orbit is inversely proportional to the azimuthal quantum number.
- (c) According to wave mechanics, the ground state angular momentum is  $h$  equal to  $\frac{h}{2\pi}$ .
- (d) The plot of  $\psi$  Vs  $r$  for various azimuthal quantum numbers, shows peak shifting towards higher  $r$  value.

(1) (b), (c) (2) (a), (d) (3) (a), (b) (4) (a), (c)





Ans. (4)

Sol. Refer Theory

8. The tests performed on compound X and their inferences are:

Test	Inference
(a) 2,4 - DNP test	Coloured precipitate
(b) Iodoform test	Yellow precipitate
(c) Azo-dye test	No dye formation


Compound 'X' is:




Ans. (2)

Sol. → 2,4 - DNP test is given by aldehyde on ketone

→ Iodoform test is given by compound having







9. The major product formed in the following reaction is:



**Complete Sample Paper of  
CBSE Class 12<sup>th</sup>  
Board Examination 2019**


**Mathematics (Solutions)  
(100% Guaranteed)**

Please Our Youtube Channel  
'Concepts Made Easy'



**Ans. (1)**

**Sol.** Aldehyde reacts at a faster rate than keton during aldol and sterically less hindered anion will be a better nucleophile so self aldol at



- 10.** For the reaction,  $2\text{A} + \text{B} \rightarrow \text{products}$ , when the concentrations of A and B both were doubled, the rate of the reaction increased from  $0.3 \text{ mol L}^{-1}\text{s}^{-1}$  to  $2.4 \text{ mol L}^{-1}\text{s}^{-1}$ . When the concentration of A alone is doubled, the rate increased from  $0.3 \text{ mol L}^{-1}\text{s}^{-1}$  to  $0.6 \text{ mol L}^{-1}\text{s}^{-1}$


Which one of the following statements is correct ?

- (1) Order of the reaction with respect to B is 2
- (2) Order of the reaction with respect to A is 2
- (3) Total order of the reaction is 4
- (4) Order of the reaction with respect to B is 1

**Ans. (1)**



**Sol.**  $r = k[\text{A}]^x[\text{B}]^y$   
 $\Rightarrow 8 = 2^3 = 2^{x+y}$   
 $\Rightarrow x + y = 3 \dots (1)$   
 $\Rightarrow 2 = 2^x$   
 $\Rightarrow x = 1, y = 2$   
 Order w.r.t. A = 1  
 Order w.r.t. B = 2

- 11.** The correct sequence of amino acids present in the tripeptide given below is :




- (1) Leu - Ser - Thr
- (2) Thr - Ser - Leu
- (3) Thr - Ser - Val
- (4) Val - Ser - Thr

**Ans. (4)**



- 12.** The correct statement regarding the given Ellingham diagram is:







## JEE (Main) Examination–2019/Evening Session/09-01-2019




- (1) At 800°C, Cu can be used for the extraction of Zn from ZnO  
 (2) At 500 C, coke can be used for the extraction of Zn from ZnO  
 (3) Coke cannot be used for the extraction of Cu from Ca<sub>2</sub>O.  
 (4) At 1400°C, Al can be used for the extraction of Zn from ZnO

**Ans. (4)****Sol.** According to the given diagram Al can reduce ZnO.

- 13.** For the following reaction, the mass of water produced from 445 g of C<sub>57</sub>H<sub>110</sub>O<sub>6</sub> is :



- (1) 495 g (2) 490 g (3) 890 g (4) 445 g

**Ans. (1)****Sol.** moles of C<sub>57</sub>H<sub>110</sub>O<sub>6</sub>(s) =  $\frac{445}{890} = 0.5$  moles

$$n_{\text{H}_2\text{O}} = \frac{110}{4} = \frac{55}{2}$$

$$m_{\text{H}_2\text{O}} = \frac{55}{2} \times 18 = 495\text{gm}$$

- 14.** The correct match between Item I and Item II is :

**Item I**

- (A) Benzaldehyde  
 (B) Alumina  
 (C) Acetonitrile

**Item II**

- (P) Mobile phase  
 (Q) Adsorbent  
 (R) Adsorbate


- (1) (A) → (Q); (B) → (R); (C) → (P)  
 (2) (A) → (P); (B) → (R); (C) → (Q)  
 (3) (A) → (Q); (B) → (P); (C) → (R)  
 (4) (A) → (R); (B) → (Q); (C) → (P)

**Ans. (4)****Sol.**

- 15.** The increasing basicity order of the following compounds is :



- (1) (D)<(C)<(A)<(B) (2) (A)<(B)<(D)<(C)  
 (3) (A)<(B)<(C)<(D) (4) (D)<(C)<(B)<(A)

**Ans. (1)****Sol.**

- 16.** For coagulation of arsenious sulphide sol, which one of the following salt solution will be most effective ?

- (1) AlCl<sub>3</sub> (2) NaCl  
 (3) BaCl<sub>2</sub> (4) Na<sub>3</sub>PO<sub>4</sub>

**Ans. (1)****Sol.** Sulphide is -ve charged colloid so cation with maximum charge will be most effective for coagulation.Al<sup>3+</sup> > Ba<sup>2+</sup> > Na<sup>+</sup> coagulating power.

- 17.** At 100°C, copper (Cu) has FCC unit cell structure with cell edge length of x Å. What is the approximate density of Cu (in g cm<sup>-3</sup>) at this temperature ?

[Atomic Mass of Cu = 63.55u]

- (1)  $\frac{105}{x^3}$  (2)  $\frac{211}{x^3}$  (3)  $\frac{205}{x^3}$  (4)  $\frac{422}{x^3}$

**Ans. (4)**




Sol. FCC unit cell  $Z = 4$

$$d = \frac{63.5 \times 4}{6 \times 10^{23} \times x^3 \times 10^{-24}} \text{ g/cm}^3$$

$$d = \frac{63.5 \times 4 \times 10}{6} \text{ g/cm}^3$$


$$d = \frac{423.33}{x^3} \approx \left( \frac{422}{x^3} \right)$$

18. The major product obtained in the following reaction is :



- (1)
- (2)
- (3)
- (4)

Ans. (3)



19. Which of the following conditions in drinking water causes methemoglobinemia ?

- (1)  $> 50$  ppm of lead  
 (2)  $> 100$  ppm of sulphate  
 (3)  $> 50$  ppm of chloride  
 (4)  $> 50$  ppm of nitrate

Ans. (4)

Sol. Concentration of nitrate  $> 50$  ppm in drinking water causes methemoglobinemia

20. Homoleptic octahedral complexes of a metal ion  $\text{M}^{3+}$  with three monodentate ligands and  $L_1, L_2, L_3$  absorb wavelengths in the region of green, blue and red respectively. The increasing order of the ligand strength is :

- (1)  $L_2 < L_1 < L_3$                       (2)  $L_3 < L_2 < L_1$   
 (3)  $L_3 < L_1 < L_2$                       (4)  $L_1 < L_2 < L_3$

Ans. (3)

Sol. Order of  $\lambda_{\text{abs}}$  -  $L_3 > L_1 > L_2$

So  $\Delta_o$  order will be  $L_2 > L_1 > L_3$  (as  $\Delta_o \propto \frac{1}{\lambda_{\text{abs}}}$ )

So order of ligand strength will be  $L_2 > L_1 > L_3$


21. The product formed in the reaction of cumene with  $\text{O}_2$  followed by treatment with dil. HCl are :

- (1) and
- (2) and  $\text{CH}_3 - \text{OH}$
- (3) and
- (4) and

Ans. (3)



## JEE (Main) Examination-2019/Evening Session/09-01-2019

**Sol. Cummene hydroperoxide reaction**

22. The temporary hardness of water is due to :-

- (1)  $Ca(HCO_3)_2$                       (2)  $NaCl$   
 (3)  $Na_2SO_4$                          (4)  $CaCl_2$


**Ans. (1)**

**Sol.**  $Ca(HCO_3)_2$  is responsible for temporary hardness of water

23. The entropy change associated with the conversion of 1 kg of ice at 273 K to water vapours at 383 K is :

- (Specific heat of water liquid and water vapour are  $4.2 \text{ kJ K}^{-1} \text{ kg}^{-1}$  and  $2.0 \text{ kJ K}^{-1} \text{ kg}^{-1}$ ; heat of liquid fusion and vapourisation of water are  $344 \text{ kJ kg}^{-1}$  and  $2491 \text{ kJ kg}^{-1}$ , respectively).  
 ( $\log 273 = 2.436$ ,  $\log 373 = 2.572$ ,  $\log 383 = 2.583$ )  
 (1)  $7.90 \text{ kJ kg}^{-1} \text{ K}^{-1}$             (2)  $2.64 \text{ kJ kg}^{-1} \text{ K}^{-1}$   
 (3)  $8.49 \text{ kJ kg}^{-1} \text{ K}^{-1}$             (4)  $4.26 \text{ kJ kg}^{-1} \text{ K}^{-1}$

**Ans. (4)**



$$\Delta S_1 = \frac{\Delta H_{\text{fusion}}}{273} = \frac{334}{273} = 1.22$$

$$\Delta S_2 = 4.2 \ln \left( \frac{363}{273} \right) = 1.31$$

$$\Delta S_3 = \frac{\Delta H_{\text{vap}}}{373} = \frac{2491}{373} = 6.67$$

$$\Delta S_4 = 2.0 \ln \left( \frac{383}{373} \right) = 0.05$$

$$\Delta S_{\text{total}} = 9.26 \text{ kJ kg}^{-1} \text{ K}^{-1}$$

24. The pH of rain water, is approximately :

- (1) 6.5            (2) 7.5            (3) 5.6            (4) 7.0

**Ans. (3)**

**Sol.** pH of rain water is approximate 5.6

25. If the standard electrode potential for a cell is 2 V at 300 K, the equilibrium constant (K) for the reaction



at 300 K is approximately.

$$(R = 8 \text{ JK}^{-1} \text{ mol}^{-1}, F = 96000 \text{ C mol}^{-1})$$

- (1)  $e^{160}$                                       (2)  $e^{320}$   
 (3)  $e^{-160}$                                     (4)  $e^{-80}$

**Ans. (1)**

**Sol.**  $\Delta G^\circ = -RT \ln k = -nFE_{\text{cell}}^\circ$

$$\ln k = \frac{n \times F \times E^\circ}{R \times T} = \frac{2 \times 96000 \times 2}{8 \times 300}$$

$$\ln k = 160$$

$$k = e^{160}$$

26. A solution containing 62 g ethylene glycol in 250 g water is cooled to  $-10^\circ\text{C}$ . If  $K_f$  for water is  $1.86 \text{ K kg mol}^{-1}$ , the amount of water (in g) separated as ice is :

- (1) 32            (2) 48            (3) 16            (4) 64

**Ans. (4)**

**Sol.**  $\Delta T_f = K_f \cdot m$

$$10 = 1.86 \times \frac{62/62}{W_{\text{kg}}}$$

$$W = 0.186 \text{ kg}$$

$$\Delta W = (250 - 186) = 64 \text{ gm}$$

27. When the first electron gain enthalpy ( $\Delta_{\text{eg}}H$ ) of oxygen is  $-141 \text{ kJ/mol}$ , its second electron gain enthalpy is :

- (1) almost the same as that of the first  
 (2) negative, but less negative than the first  
 (3) a positive value  
 (4) a more negative value than the first


**Ans. (3)**

**Sol.** Second electron gain enthalpy is always positive for every element.








28. The major product of the following reaction is :



Ans. (3)



29. Which of the following compounds is not aromatic ?




Ans. (3)

**Concepts Made Easy**  
**JEE Mains & Advanced**  
**Free Online Exam**  
**Give test & Check your All India Rank**  
[www.ConceptsMadeEasy.com](http://www.ConceptsMadeEasy.com)

Click Here for JEE Mains & Advanced Free Online Test Practice

<https://examination-system.shineitsolutions.in/jee-mains-and-advanced-examination-system/student-login.php>


<https://examination-system.shineitsolutions.in/jee-mains-and-advanced-examination-system/student-login.php>



Do not have  $(4n + 2)$   $\pi$  electron It has  $4n$   $\pi$  electrons

So it is Anti aromatic.

30. Consider the following reversible chemical reactions :





The relation between  $K_1$  and  $K_2$  is :

(1)  $K_2 = K_1^3$  (2)  $K_2 = K_1^{-3}$

(3)  $K_1K_2 = 3$  (4)  $K_1K_2 = \frac{1}{3}$

Ans. (2)



$$\Rightarrow \left(\frac{1}{K_1}\right)^3 = K_2 \Rightarrow K_2 = (K_1)^{-3}$$

**Concepts Made Easy**  
**CBSE NEET & AIIMS Exam 2019**  
**Free Online Exam**  
**Give test & Check your All India Rank**  
[www.ConceptsMadeEasy.com](http://www.ConceptsMadeEasy.com)

Click Here for CBSE NEET & AIIMS Free Online Test Practice

<https://examination-system.shineitsolutions.in/cbse-neet-and-aiims-examination-system/student-login.php>

<https://examination-system.shineitsolutions.in/cbse-neet-and-aiims-examination-system/student-login.php>